

Leptoproduction of J/ψ

Thomas Mehen

California Institute of Technology, Pasadena, CA 91125
E-mail: mehen@theory.caltech.edu

I review the status of the extraction of NRQCD color-octet J/ψ production matrix elements. Large theoretical uncertainties in current extractions from hadroproduction and photoproduction are emphasized. Leptoproduction of J/ψ is calculated within the NRQCD factorization formalism. Color-octet contributions dominate the cross section, allowing for a reliable extraction of $\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle$ and $\langle \mathcal{O}_8^{J/\psi}(^3P_0) \rangle$. A comparison with preliminary data from the H1 collaboration shows that leading order color-octet mechanisms agree with the measured total cross section for $Q^2 > 4m_c^2$, while the color-singlet model underpredicts the cross section by a factor of 5.

The NRQCD factorization formalism of Bodwin, Braaten, and Lepage¹ has emerged as a new paradigm for computing the production and decay rates of heavy quarkonia. This formalism provides a rigorous theoretical framework which systematically incorporates relativistic corrections and ensures the infrared safety of perturbative calculations.² In the NRQCD factorization formalism, cross sections for the production of a quarkonium state H are written as

$$\sigma(H) = \sum_n \frac{c_n(\alpha_s, m_Q)}{m_Q^{d_n-4}} \langle 0 | \mathcal{O}_n^H | 0 \rangle, \quad (1)$$

where m_Q is the mass of the heavy quark Q . The short-distance coefficients, c_n , are associated with the production, at distances of order $1/m_Q$ or less, of a $Q\bar{Q}$ pair with quantum numbers indexed by n (angular momentum, $^{2S+1}L_J$, and color, 1 or 8). They are computable in perturbation theory. In Eq. (1), $\langle 0 | \mathcal{O}_n^H | 0 \rangle$ are vacuum matrix elements of NRQCD operators:

$$\langle 0 | \mathcal{O}_n^H | 0 \rangle \equiv \sum_X \sum_\lambda \langle 0 | \mathcal{K}_n^\dagger | H(\lambda) + X \rangle \langle H(\lambda) + X | \mathcal{K}_n | 0 \rangle, \quad (2)$$

where $\mathcal{K}_n$ is a bilinear in heavy quark fields which creates a $Q\bar{Q}$ pair in an angular-momentum and color configuration indexed by n . The bilinear combination $\mathcal{K}_n^\dagger \mathcal{K}_n$ has energy dimension d_n . The production matrix elements describe the evolution of the $Q\bar{Q}$ pair into a final state containing the quarkonium H plus additional hadrons (X) which are soft in the quarkonium rest frame. Throughout the remainder of this talk, a shorthand notation will be used in which the vacuum matrix elements are written as $\langle \mathcal{O}_{(1,8)}^H(^{2S+1}L_J) \rangle$.

The NRQCD matrix elements obey simple scaling laws³ with respect to v , the relative velocity of the Q and $\bar{Q}$ in the quarkonium bound state. Therefore, Eq. (1) is a double expansion in v and α_s . Since the NRQCD matrix elements are sensitive only to large distance scales, they are independent of the short-distance process in which the Q and $\bar{Q}$ are produced. Thus, the NRQCD matrix

elements are universal parameters which can be extracted from one experiment and used to predict production cross sections in other processes.

Prior to the innovations presented in Ref. ¹, most J/ψ production calculations took into account only the hadronization of $c\bar{c}$ pairs initially produced in a color-singlet 3S_1 state, as parameterized by the NRQCD matrix element $\langle \mathcal{O}_1^{J/\psi}(^3S_1) \rangle$. An important aspect of the NRQCD formalism is that, in addition to the color-singlet contribution, it allows for the possibility that a $c\bar{c}$ pair produced in a color-octet state can evolve non-perturbatively into a J/ψ . The most important color-octet matrix elements are $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$, $\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle$, and $\langle \mathcal{O}_8^{J/\psi}(^3P_J) \rangle$, which are suppressed by v^4 relative to the leading color-singlet matrix element. They describe the non-perturbative evolution of a color-octet $c\bar{c}$ pair in either a 3S_1 , 1S_0 , or 3P_J angular momentum state into a J/ψ . Using heavy quark spin symmetry relations, it is possible to express all three P-wave matrix elements in terms of one: $\langle \mathcal{O}_8(^3P_J) \rangle = (2J+1) \langle \mathcal{O}_8(^3P_0) \rangle + O(v^2)$. Thus, at this order in the v expansion, there are three independent color-octet matrix elements.

The NRQCD factorization formalism has enjoyed considerable phenomenological success. Most notably, the large $p_\perp$ production of J/ψ at hadron colliders, which is underpredicted by over an order of magnitude in the color-singlet model (CSM), can be easily accounted for by color-octet production mechanisms.⁴ The CSM cannot account for the observed branching ratios for $B \rightarrow J/\psi + X$, $B \rightarrow \psi' + X$. Inclusion of color-octet mechanisms removes this discrepancy.^{5,6} Color-octet mechanisms also improve the understanding of J/ψ production in Z^0 decay.⁷

The NRQCD factorization formalism has yet to be conclusively proven as the correct theory of quarkonium production. One important test of the formalism is the verification of the universality of the NRQCD matrix elements. Another important prediction, which has yet to be verified, is the polarization of J/ψ at large $p_\perp$ at

hadron colliders. Large $p_\perp$ production of J/ψ is dominated by fragmentation of a (nearly) on-shell gluon into a $^3S_1^{(8)} c\bar{c}$ pair, which inherits the gluon's transverse polarization. Because of heavy quark spin symmetry, soft gluons emitted as the $c\bar{c}$ pair hadronizes into the quarkonium state do not dilute this polarization.^{8,9} Therefore, at large $p_\perp$, J/ψ are expected to be almost 100% transversely polarized. This gets significantly diluted at lower $p_\perp$ due to nonfragmentation production.^{10,11}

Polarization is a particularly interesting test of quarkonium production because it can distinguish between NRQCD and the other remaining model of quarkonium production, the color evaporation model (CEM).¹² In this model, the cross section for producing a J/ψ is proportional to the total production rate for $c\bar{c}$ pairs with invariant mass less than the $D\bar{D}$ threshold:

$$\sigma_{J/\psi} = \rho_{J/\psi} \int_{2m_c}^{2m_D} dm_{c\bar{c}} \frac{d\sigma_{c\bar{c}}}{dm_{c\bar{c}}}, \quad (3)$$

where $\rho_{J/\psi}$ is a universal factor which represents the fraction of $c\bar{c}$ pairs which hadronize into a J/ψ . Because this model includes color-octet production mechanisms, it is consistent with hadron collider data as well as data from Z^0 decay¹². A prediction of this model is that the emission of soft gluons in the hadronization washes out any polarization of the $c\bar{c}$ produced in the short distance process. This prediction is at odds with heavy quark spin symmetry arguments. The CEM would be a reasonable model if the relative velocity, v , in the quarkonium state were too large to serve as a useful expansion parameter. From the existing estimates of the color-octet matrix elements, this does not appear to be the case. The leading color-singlet matrix element is well determined: $\langle \mathcal{O}_1^{J/\psi}(^3S_1) \rangle = 1.1 \pm 0.1 / \text{GeV}^3$. Extractions of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$, $\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle$, and $\langle \mathcal{O}_8^{J/\psi}(^3P_0) \rangle / m_c^2$ suffer from considerable uncertainty but they are known to be of order 10^{-2} GeV^3 . These values are consistent with the v^4 suppression expected on the basis of NRQCD scaling laws.

To quantitatively predict polarization as a function of $p_\perp$, color-octet matrix elements need to be accurately determined. At the present time, extractions from production at hadron colliders suffer from considerable theoretical uncertainty. Beneke and Krämer¹¹ extract $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle = 1.2_{-0.7}^{+1.2} \times 10^{-2} \text{ GeV}^3$, where statistical errors, and errors due to variation of the renormalization scale and the parton distribution functions, have been added in quadrature. The error is dominated by scale uncertainty. Other extractions^{13,14} of $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$ are consistent within these errors. $\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle$, $\langle \mathcal{O}_8^{J/\psi}(^3P_0) \rangle$ are even more poorly determined since hadroproduction is only sensitive to a particular linear combination. The extracted value of this linear combination is also much

more sensitive to the choice of parton distribution function and the effects of initial and final state radiation.^{15,16} Extractions from lower energy hadroproduction¹⁷ suffer similar from problems. These calculations are extremely sensitive to the charm quark mass because of the small energy scale of the process. Corrections due to higher orders in perturbation theory and higher twist effects are expected to be large.

Recently, progress has been made in analysis of B decays and Z^0 decay. A next-to-leading order analysis of B decay⁶ establishes a bound on the following linear combination of color-octet matrix elements:

$$\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle + 3.1 \frac{\langle \mathcal{O}_8^{J/\psi}(^3P_0) \rangle}{m_c^2} < 2.3 \times 10^{-2} \text{ GeV}^3. \quad (4)$$

The individual matrix elements, $\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle$ and $\langle \mathcal{O}_8^{J/\psi}(^3P_0) \rangle$ remain undetermined. Production of J/ψ in Z^0 decays is dominated by gluon fragmentation and is therefore sensitive to $\langle \mathcal{O}_8^{J/\psi}(^3S_1) \rangle$. A recent analysis,¹⁸ which sums logarithms of $2M_{J/\psi}/M_Z$ and $2E_{J/\psi}/M_Z$ to all orders, extracts the following effective matrix element:

$$\begin{aligned} \langle \hat{\mathcal{O}}_8^{J/\psi}(^3S_1) \rangle &= \sum_H \langle \hat{\mathcal{O}}_8^H(^3S_1) \rangle \text{Br}(H \rightarrow J/\psi + X) \\ &= 1.9 \pm 0.5 \pm 1.0 \times 10^{-2} \text{ GeV}^3. \end{aligned} \quad (5)$$

The first error is statistical, the second theoretical. The effective $\langle \hat{\mathcal{O}}_8^{J/\psi}(^3S_1) \rangle$ includes feeddown from higher mass charmonium resonances (H) because it is not possible to separate prompt J/ψ experimentally.

Next-to-leading order calculations of photoproduction would seem to indicate that NRQCD predicts a large excess of J/ψ at large z which is not observed in the data.¹⁹ (Here $z = E_{J/\psi}/E_\gamma$, where energies are measured in the proton rest frame.) This led to the conclusion that the color-octet matrix elements extracted from the Tevatron were an order of magnitude too large to be consistent with photoproduction data from the HERA experiments. However, this enhancement is actually an artifact of the next-to-leading order perturbative calculation. It is worth examining this process in more detail because it is indicative of some of the pitfalls one may encounter when trying to compare NRQCD with data.

The leading order diagrams are shown in Fig. 1, and results in the following perturbative cross section:

$$\frac{d\sigma}{dz} = \sigma_0 \delta(1-z), \quad (6)$$

$$\begin{aligned} \sigma_0 &= \frac{4\pi^3 \alpha_s \alpha}{s_{\gamma p} m_c^3} G(4m_c^2/s_{\gamma p}) \times \\ &\quad \left(\langle \mathcal{O}_8^{J/\psi}(^1S_0) \rangle + 7 \frac{\langle \mathcal{O}_8^{J/\psi}(^3P_0) \rangle}{m_c^2} \right) \end{aligned} \quad (7)$$




Figure 1: Leading order diagrams for color-octet photoproduction of J/ψ

$G(x_g)$ is the gluon distribution function of the proton. $s_{\gamma p}$ is the center of mass energy squared of the photon-proton system. The delta function in Eq.(6) is an artifact of the perturbative expansion which will be smeared by nonperturbative emission of soft gluons as the $c\bar{c}$ pair hadronizes into the final state hadron. How this nonperturbative smearing can be accounted for within the NRQCD formalism will be discussed below. For now, we concern ourselves with perturbative corrections.

The next-to-leading order diagrams are shown in Fig. 2. In the limit in which the final state gluon is soft, the cross section is given by: ²⁰

$$\begin{aligned} \frac{d\sigma}{dz} &= \int dx_g G(x_g) \frac{\sum |M_{LO}|^2}{16\pi x_g s_{\gamma p}} \\ &\quad \times C_A g_s^2 \left(\frac{2P \cdot g}{P \cdot k g \cdot k} - \frac{4m_c^2}{(P \cdot k)^2} \right) \\ &= \int_\rho^z dx G(\rho/x) \frac{\sum |M_{LO}|^2}{4m_c^2 s_{\gamma p}} C_A \alpha_s \frac{z-x}{(1-x)^2(1-z)} \\ &= -\sigma_0 \frac{C_A \alpha_s}{\pi} \frac{\ln(1-z)}{1-z} + \dots \end{aligned} \quad (8)$$

M_{LO} is the leading order matrix element from Fig. 1. P , g , and k are the four momentum of the J/ψ , the initial state gluon, and the final state gluon, respectively. $\rho = 4m_c^2/s_{\gamma p}$ and $x = \rho/x_g$. In the last line of Eq.(8), we take $x = 1$ inside the argument of the structure function in order to obtain an analytic expression. This approximation is valid up to subleading logs since the leading logs come from the region $x \rightarrow z \rightarrow 1$. In higher orders, we expect to find terms of the form $\alpha_s^n \ln^{2n-1}(1-z)/(1-z)$ which need to be resummed before theory can be compared with experimental data. The divergence of the cross section as $z \rightarrow 1$ is an artifact of the next-to-leading order calculation. Once the resummation is performed, the cross section will be well-behaved as $z \rightarrow 1$.

There are also important nonperturbative corrections²¹ near $z = 1$. For quarkonium production near the boundaries of phase space, it is sometimes the case that contributions from NRQCD operators which




Figure 2: Next-to-leading order diagrams for color-octet photoproduction of J/ψ

are higher order in v are enhanced by kinematic factors. This results in the breakdown of the NRQCD expansion. The crux of the problem is that in the perturbative QCD part of the matching calculation one uses twice the heavy quark mass instead of the quarkonium mass to compute the phase space for the production of the quarkonium meson. The difference between $2m_Q$ and M_H is a v^2 correction, which is ignored in leading-order calculations. However, at the boundaries of phase space this difference becomes important, and it is necessary to sum an infinite number of NRQCD matrix elements. This resummation leads to a universal distribution function called a shape function which replaces the delta function in Eq. (6). Because of the universality of the shape functions, it may be possible in the future to test NRQCD by comparing shape functions extracted from different quarkonium production processes. This would require more precise data than is currently available.

As $z \rightarrow 1$, the photoproduction cross section becomes sensitive to the effects of soft gluon radiation. Within perturbation theory this sensitivity is signalled by the presence of large logarithms which must be resummed. The v expansion in Eq. (1) also breaks down and an infinite set of operators become relevant. For this reason, it is impossible to extract NRQCD matrix elements from this distribution. When leptonproduction is discussed below, the analysis will focus on observables for which shape function corrections are not kinematically enhanced, because it is only from these distributions where one can hope to obtain a reliable extraction of NRQCD matrix elements.

In Ref.²², leptonproduction of J/ψ is examined and found to be a useful process from which to extract NRQCD matrix elements. This process is similar to photoproduction, however, now the photon is off-shell. At leading order, the cross section is again given by the diagrams in Fig. 1:

$$\frac{d\sigma}{dQ^2} = \int dy \int dx_g G(x_g) \delta(x_g y s - (2m_c)^2 - Q^2)$$

$$\times \frac{2\alpha_s(\mu)\alpha^2 e_c^2 \pi^2}{Q^2(Q^2 + (2m_c)^2)m_c} \left\{ \frac{1 + (1-y)^2}{y} \left[\langle \mathcal{O}_8^\psi(^1S_0) \rangle + \frac{3Q^2 + 7(2m_c)^2}{Q^2 + (2m_c)^2} \frac{\langle \mathcal{O}_8^\psi(^3P_0) \rangle}{m_c^2} \right] - \frac{8(2m_c)^2 Q^2}{x_g s(Q^2 + (2m_c)^2)} \frac{\langle \mathcal{O}_8^\psi(^3P_0) \rangle}{m_c^2} \right\}, \quad (9)$$

where s is the electron-proton center-of-mass energy squared. The momentum fraction of the virtual photon relative to the incoming lepton is $y \equiv P_p \cdot q / P_p \cdot k$, where P_p is the proton four-momentum, q is the photon four-momentum, and k is the incoming lepton four-momentum, and $Q^2 \equiv -q^2$.

There are several advantages to extracting NRQCD matrix elements from this process. First of all, note that the relative importance of $\langle \mathcal{O}_8^\psi(^1S_0) \rangle$ and $\langle \mathcal{O}_8^\psi(^3P_0) \rangle$ changes as a function of Q^2 . Thus, it is possible to fit the differential cross section as a function of Q^2 and extract both of these matrix elements separately. Second, the scale for the coupling is set by $\mu^2 \approx Q^2 + (2m_c)^2$, so as Q^2 increases, perturbative corrections should be increasingly suppressed relative to photoproduction. In Ref.²², next-to-leading order graphs with real gluon emission were computed and found to give a small contribution to the total cross section for $Q^2 > 4 \text{ GeV}^2$. The uncertainty due to varying the renormalization scale is less than 10% for $Q^2 > 4 \text{ GeV}^2$. This is a great deal smaller than the corresponding uncertainty in hadroproduction. This indicates that higher order perturbative corrections to the leading order color-octet production mechanism will be small. Higher twist corrections to the parton model are also expected to be small as they are suppressed by powers of Q^2 .

The leading color-singlet mechanism requires the emission of an additional hard gluon. (See the first diagram in Fig. 2.) This $\alpha_s(Q)$ suppression compensates for the v^4 suppression of the color-octet matrix elements, and it turns out that for $Q^2 > 4 \text{ GeV}^2$, the color-octet contribution is roughly 4 times bigger than the leading color-singlet process. However, color-singlet mechanisms will dominate in the inelastic ($z < 1$) region, where there is no $O(\alpha_s)$ color-octet contribution. In the elastic region, one also expects production of J/ψ via diffractive processes. At large Q^2 , diffractive leptonproduction can be studied using perturbative QCD. In Ref.²³, a perturbative analysis of diffractive leptonproduction predicts the cross section to fall as $1/(Q^2 + (2m_c)^2)^3$, as compared to $1/(Q^2 + (2m_c)^2)^2$ for the color-octet mechanism. Therefore, at sufficiently large Q^2 , the diffractive contribution should be negligible correction to our calculation.

One must also consider the possibility of kinematic enhancement of higher order v^2 corrections. An analysis of the Q^2 distribution in Ref.²² shows that the higher

order in v^2 corrections associated with the shape function are suppressed in the large Q^2 limit. Therefore, this distribution is calculable in NRQCD. However, the z distribution still suffers from large endpoint corrections, and a perturbative calculation of this distribution cannot be compared with experiment near $z = 1$, even at large Q^2 .

The largest errors in the calculation of the Q^2 distribution are associated with uncertainty in the charm quark mass. Varying m_c between 1.3 GeV and 1.7 GeV, results in an error of $^{+60\%}_{-25\%}$ at $Q^2 = 10 \text{ GeV}^2$. This error decreases slightly as Q^2 is increased.

Once color-octet matrix elements are extracted, the polarization of the J/ψ can be predicted without introducing any new parameters. Calculations of the polarization of J/ψ produced via the leading order color-octet mechanism as a function of Q^2 are given in Ref.²². A precise measurement of the polarization of leptonproduced J/ψ could provide an excellent test of the NRQCD factorization formalism.

Preliminary results from HERA,²⁴ indicate that $d\sigma/dQ^2$ is well described by NRQCD for $Q^2 > (2m_c)^2$. This is in contrast with CSM calculations which underpredict the cross section by a factor of ~ 5 . So this appears to be strong evidence in favor of color-octet mechanisms as understood within the NRQCD factorization formalism. However, the shape of the rapidity distribution at large rapidity is not well reproduced by NRQCD. This could be a consequence of the effects of soft gluon emission which are not accounted for in the leading order calculations of Ref.²². Analysis of soft gluon effects and extraction of NRQCD matrix elements will be performed in a future publication.²⁵

Acknowledgements

This work was done in collaboration with Sean Fleming. This work was supported by the National Science Foundation under grant number PHY-94004057, by the Department of Energy under grant number DE-FG03-92-ER40701, and by a John A. McCone Fellowship.

References

1. G.T. Bodwin, E. Braaten, and G.P. Lepage, Phys. Rev. D **51** 1125 (1995).
2. H.W. Huang and K.T. Chao, Phys. Rev. D **55** 244, (1997); E. Braaten and Y.Q. Chen, Report No. OHSTY-HEP-T-97-001, hep-ph/9701242 (unpublished).
3. G.P. Lepage, L. Magnea, C. Nakhleh, U. Magnea, and K. Hornbostle, Phys. Rev. D **46**, 4052 (1992).
4. E. Braaten and S. Fleming, Phys. Rev. Lett. **74**3327 (1995).

5. P. Ko, J. Lee, and H.S. Song, Phys. Rev. **D53**, 1409 (1996).
6. M. Beneke, F. Maltoni, and I. Z. Rothstein, hep-ph/9808360.
7. P. Cho, Phys. Lett. **B368** 171 (1996).
8. P. Cho and M. Wise, Phys. Lett. **B346** 129 (1995).
9. M. Beneke and I. Z. Rothstein, Phys. Lett. **B372** 157 (1996), **B389** 769 (1996).
10. A. Leibovich, Phys. Rev. **D56** 4412 (1997).
11. M. Beneke and M. Krämer, Phys. Rev. **D55**, 5269 (1997).
12. J.F. Amundson, O.J.P. Éboli, E. M. Gregores, and F. Halzen, Phys. Lett. B **372**, 127 (1996), *ibid* **390** 323 (1997); O.J.P. Éboli, E. M. Gregores, and F. Halzen, Phys. Lett. B **395**, 113 (1997); G. A. Schuler and R. Vogt, Phys. Lett. B **387**, 181 (1996).
13. P. Cho and A. Leibovich, Phys. Rev. D **53**, 150 (1996); **53**, 6203 (1996).
14. M. Cacciari, M. Greco, M.L. Mangano, and A. Petrelli, Phys. Lett. **356B**, 553 (1995).
15. B. A. Kniehl and G. Kramer, hep-ph/9803256.
16. B. Cano-Coloma and M.A. Sanchis-Lozano, Nucl. Phys. **B508**, 753 (1997); Phys. Lett. **B406**, 232 (1997).
17. M. Beneke and I. Z. Rothstein, Phys. Rev. **D54** 2005 (1996), **D54** 7082 (1996).
18. C.G. Boyd, A. Leibovich, and I. Rothstein, hep-ph/9810364.
19. M. Cacciari and M. Krämer, Phys. Rev. Lett. **76**, 4128 (1996); P. Ko, J. Lee, and H.S. Song, Phys. Rev. **D54**, 4312 (1996).
20. F. Maltoni, M.L. Mangano, and A. Petrelli, Nucl. Phys. **B519**, 361 (1998).
21. M. Beneke, I. Z. Rothstein, and M. B. Wise, Phys. Lett. **B408**, 373 (1997).
22. S. Fleming and T. Mehen, Phys. Rev. **D57**, 1846 (1998).
23. S. J. Brodsky, L. Frankfurt, J. F. Gunion, A.H. Mueller, and M. Strikman, Phys. Rev. D **50**, 3134 (1994);
M.G. Ryskin, R.G. Roberts, A.D. Martin, and E.M. Levin, Report No. RAL-TR-95-065, hep-ph/9511228.
24. H1 Collaboration, preprint submitted to ICHEP 98.
25. S. Fleming and T. Mehen, in preparation.